

**The effect of structured patient education on physical activity in patients with peripheral arterial disease and intermittent claudication: a systematic review**

Abaraogu, U.O.; Dall, P.M.; Seenan, C.A.

*Published in:*  
European Journal of Vascular and Endovascular Surgery

*DOI:*  
[10.1016/j.ejvs.2017.04.003](https://doi.org/10.1016/j.ejvs.2017.04.003)

*Publication date:*  
2017

*Document Version*  
Author accepted manuscript

[Link to publication in ResearchOnline](#)

*Citation for published version (Harvard):*  
Abaraogu, UO, Dall, PM & Seenan, CA 2017, 'The effect of structured patient education on physical activity in patients with peripheral arterial disease and intermittent claudication: a systematic review', *European Journal of Vascular and Endovascular Surgery*, vol. 54, no. 1, pp. 58-68. <https://doi.org/10.1016/j.ejvs.2017.04.003>

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.

**Title:** The effect of structured patient education on physical activity in patients with peripheral arterial disease and intermittent claudication: a systematic review.

**Short title:** patient education in intermittent claudication

**Category:** Review

**Authors:** <sup>a,b</sup>Ukachukwu O Abaraogu, BMR, MSc PT; <sup>a</sup>Philippa M Dall, PhD; <sup>a</sup>Christopher Seenan, PhD PT.

**Institutional Affiliations:** <sup>a</sup>School of Health and Life Sciences, Glasgow Caledonian University, Glasgow, United Kingdom. <sup>b</sup>Department of Medical Rehabilitation, University of Nigeria Enugu Campus Enugu, Nigeria.

**Corresponding author:** Ukachukwu O Abaraogu | School of Health and Life Sciences, Glasgow Caledonian University, Glasgow, United Kingdom Email:

[Ukachukwu.Abaraogu@gcu.ac.uk](mailto:Ukachukwu.Abaraogu@gcu.ac.uk); Department of Medical Rehabilitation University of Nigeria Enugu Campus Enugu 400006 Enugu Nigeria. Email:

[Ukachukwu.Abaraogu@unn.edu.ng](mailto:Ukachukwu.Abaraogu@unn.edu.ng)

20 **What does this review add to the existing literature and how will it influence future**  
21 **clinical practice:** This article describes the first comprehensive systematic review of patient  
22 education interventions for physical activity (PA) improvement for individuals with  
23 intermittent claudication. The current evidence is inconclusive regarding the effect of  
24 patients' education on PA of individuals with IC. Further research is warranted to establish  
25 the effects and optimal design of education interventions.

26

27 **Manuscript word count** (Including title page, review highlights, abstract text, and  
28 references): 4516

29

30

## 31 **Abstract**

32 **Objectives:** To review the components and effects of patient education interventions to  
33 improve PA in patients with PAD and IC, and patients' experiences of these interventions.

34 **Data Sources:** CINAHL, Cochrane Library, Ovid, ProQuest, AMED, MEDLINE,  
35 PsycINFO, *Web of Science Core Collection*, and PEDRO, and Trial registers and directory of  
36 Open-Access repository websites and Web of science conference proceedings were searched.  
37 Hand searching of reference lists of identified studies was also performed to identify studies  
38 that reported the effect of patient education interventions on daily PA and/or walking  
39 capacity in individuals with PAD and IC, or studies investigating patients' experiences of  
40 such interventions.

41 **Review Methods:** A systematic search was conducted in June 2016 (updated in March  
42 2017). Primary outcomes were daily step count and self-reported PA; the secondary outcome  
43 was absolute claudication distance. There was substantial heterogeneity in terms of  
44 modalities of patient education in the included studies; hence a narrative synthesis  
45 was implemented.

46 **Results:** Six studies (1087 participants) were included in the review. Findings from a small  
47 number of high-quality trials demonstrated potential for PA improvement with structured  
48 education interventions. Nevertheless, evidence is currently inconclusive regarding the effect  
49 on daily PA and walking capacity of patients with IC. Patients reported that they valued the  
50 interventions studied, finding them acceptable and important in improving their PA,  
51 motivating and empowering them to self-manage their condition.

52 **Conclusions:** The evidence from the review is limited and inconclusive regarding the  
53 effectiveness of structured education for increasing PA in patients with PAD and IC. More  
54 rigorous trials are needed before recommendations can be made. Future interventions should

55 consider the key criteria for a structured patient education programme, and also report  
56 patients' experiences and perceptions.

57 **Abstract word count:** 274

58 **Key words:** Peripheral arterial disease, intermittent claudication, physical activity, patient  
59 education, systematic review.

60

61

## Introduction

Peripheral arterial disease (PAD) is a growing public health burden. PAD leads to arterial stenosis and consequently inadequate blood flow to the peripheries.<sup>1,2</sup> This commonly presents as pain in the lower limb(s) precipitated by exercise and relieved by rest,<sup>3</sup> defined as intermittent claudication (IC). Patients with IC experience functional decline and limitation in physical activity (PA),<sup>4</sup> further raising the risk of a cardiovascular event in a vascular system already compromised by the underlying atherosclerosis.<sup>1,3</sup> Therefore, patients with PAD present with an increased risk of cardiovascular and cerebrovascular events similar to those with coronary heart disease.<sup>5</sup>

Lower daily PA levels have been recognised as a strong predictor of increased morbidity and mortality in this population.<sup>6</sup> Supervised exercise programmes (SEPs) are recommended as a primary therapy for this population,<sup>7</sup> and have been shown to improve treadmill walking distances of patients with IC.<sup>8,9</sup> However, most studies reporting such improvements did not investigate daily PA, and those that have, did not find improvements in daily PA.<sup>9</sup> Reduced self-efficacy, attributed to poor understanding of the disease and uncertainty regarding the importance of walking has been identified as a major barrier to exercise uptake in this population.<sup>10,11</sup> These findings suggest that a patient-centred self-management approach to improving PA, including structured patient education, may be beneficial in this population.

Although current literature supports educating patients with IC about their disease pathology and the importance of walking,<sup>10,11</sup> neither evidence of effectiveness nor patients' perceptions of interventions have been established. The aim of this review is to examine the effect of patient education for improving PA in individuals with PAD and IC and the experiences and perceptions of patients of these interventions.

## **Methods**

The protocol for this review was registered with the International Prospective Register of Systematic Reviews (CRD42015027314), and has been published elsewhere.<sup>12</sup>

### ***Eligibility Criteria***

Studies reporting the effect of patient education interventions on daily PA and/or walking capacity in individuals with PAD and IC, or studies investigating patients' experiences of such interventions were included in this review. Diagnosis of PAD could be objective (eg, an ankle-brachial index (ABI) <0.9), by questionnaire or clinical diagnosis. Original English language research manuscripts in peer review journals and conference proceedings were included. Studies were included only if they reported on structured patient education interventions and/or components particularly related to PAD and IC. For the purposes of this review, the key criteria used to define structured education in diabetes were adopted.<sup>13</sup> To be included, a patient education intervention should: i) aim to empower and inform patients, and to support self-management of their PAD/IC by building sufficient knowledge and skills to do so; ii) include topics about the nature of PAD/IC, and day-to-day living and management of PAD/IC including importance of physical activity and walking; iii) have embedded quality assurance processes including having a structured curriculum, having trained educators, being quality assured, and being audited.

### ***Outcomes***

Daily PA (daily step count and self-reported change in daily PA) was the primary outcomes. Secondary outcomes included treadmill measured walking capacity (absolute claudication distance (ACD)), pain intensity, quality of life, and qualitative data regarding patients' experiences with interventions.

### ***Information sources, search strategy, study records and data management***

A systematic search was conducted in June 2016 (Updated in March 2017). Nine databases (CINAHL, Cochrane Library, Ovid, ProQuest, AMED, MEDLINE, PsycINFO, Web of Science Core Collection, and PEDRO), trial registers and directory of Open-Access repository websites were searched by the first author (UOA) using key words: patient education, lifestyle education, behaviour change intervention, peripheral arterial disease, intermittent claudication, physical activity and homebased exercise combined with specific search terms and strategies for each database.<sup>12</sup> Reference lists of identified studies were also searched. Titles, abstracts and full text of selected studies were independently screened by two authors from a pool of three (UOA, CS, PD) using previously defined eligibility criteria.<sup>12</sup> Differences of opinion regarding inclusion or exclusion were resolved by discussion between authors and reflection in consultation with the second author (PD).

### ***Data collection Processes***

The Cochrane Consumers and Communication Review Group Data Extraction Template,<sup>14</sup> and the Supplementary Guidance for Inclusion of Qualitative Research<sup>15</sup> were adapted to extract data from the included studies. The Cochrane Collaboration risk of bias tool was used to determine and summarise risk of the included studies.<sup>16</sup> Assessment was made in each of the included studies and graded as ‘high risk’ or ‘low risk’ following a well described procedure.<sup>17</sup> Studies were subsequently rated as low-quality trials (i.e. having high risk of bias) or high-quality’ trials (i.e. having low to moderate risk of bias) if there were  $\geq 3$  or  $< 3$  identifiable sources of bias respectively (Table 2). Two reviewers (UOA, CS) performed the data extraction, and made judgements regarding the risk of bias independent of each other independently. Again any disagreement was resolved by discussion between the reviewers and consultation with the second author (PD).

### **Results**



## ***Study inclusion***

The search initially identified a total of 5707 studies (Figure 1), of which six studies, contributing data on 1087 participants were included in the final analysis. Meta-analysis was not possible due to wide variations in interventions and substantial methodological and clinical heterogeneity in the included studies. The results of this review are reported using narrative synthesis.

## ***Characteristics of included studies***

***Study design, participants and quality appraisal:*** Five of the included studies were RCTs,<sup>18-22</sup> and one was a pretest-posttest design.<sup>32</sup> The number of participants in each study ranged from 23<sup>19</sup> to 882<sup>20</sup>. The basis for IC diagnosis in most (n=4) studies was post exercise ABI $\leq$ 0.9.<sup>18,20,22,30</sup> Clinical characteristics between the intervention and control groups were similar at baseline for all included studies, except for one study<sup>22</sup> where the control group had a higher resting heart rate and weight (Table 1). Overall, four of the six trials were rated as high-quality<sup>18-20,22</sup> (Table 2). No study was assessed to have a risk of bias related to selective reporting. Sources of bias in included studies included lack of blinding of outcome assessment (n=5),<sup>18-21</sup> lack of allocation concealment (n=3),<sup>18,19,23</sup> lack of participants and/or personnel blinding (n=3),<sup>21-23</sup> and not being powered to detect an effect size (n=4).<sup>18,19,21,22</sup>

## ***Components of interventions in included studies***

The included studies had wide variation in the intervention components used, but all included an education session, exercise prescription, and some behavioural change techniques (BCTs) (Table 1). Information provision, goal setting, action planning and feedback were reported in all included studies. Other reported BCTs included motivational interviewing (n=3), barrier identification/problem solving (n=5), feedback on performance (n=2), and prompting self-monitoring of behavioural outcome (n=4). Intervention duration ranged from 6 weeks<sup>19</sup> to 4 months.<sup>18</sup> Most studies (n=4) instructed patients to walk through moderate-to-severe leg pain.

Control participants either received usual care,<sup>18,19,22</sup> an active control (7-minute video)<sup>31</sup> or no intervention.<sup>20</sup> One study did not include a control group.<sup>23</sup>

### ***Outcomes reported in included studies***

Five studies<sup>18-20,22,30</sup> reported walking capacity outcomes, and three<sup>18,19,21</sup> reported outcomes of daily PA. Walking capacity outcomes included treadmill walking distances and time, six minutes walking distance, WELCH score, walking impairment questionnaire speed, distance and stair climbing scores. Daily PA was assessed objectively in two studies: daily step counts using a pedometer<sup>18</sup> and accelerometer.<sup>19</sup> Three studies<sup>18,19,23</sup> reported outcomes on patients' experiences to interventions. Other outcomes reported included pain intensity,<sup>23</sup> self-efficacy and self-esteem,<sup>19</sup> and quality of life.<sup>19-20,22,23</sup>

### **Effect of interventions in included studies**

Except where specified otherwise, effects of intervention are reported as comparison of the intervention versus the control.

### ***Daily physical activity outcomes***

***Self-report daily physical activity outcome.*** Two studies provided data on self-report of daily PA.<sup>20,21</sup> A high-quality trial,<sup>20</sup> showed no difference between percentage of participants who engaged in walking for recreation  $\geq 3$  times per week (36.8 vs 31.4%;  $p=0.14$ ), engaged in more than usual activity (10.2 vs 9.0%;  $p=0.73$ ), or belonged to an exercise group (14.0 vs 10.1%) at 2-months. However, at 12 months follow up, a greater percentage were walking  $\geq 3$  times/week (33.8 vs 25.0%;  $p=0.01$ ), engaged in more activity than usual (11.1 vs 5.9%;  $p=0.03$ ), or belonged to an exercise group (16.5 vs 1.8%;  $p<0.001$ ). The other trial was of low-quality<sup>31</sup> and reported no change in time spent in various levels of activity.

***Objectively measured daily physical activity outcome.*** Data on daily step counts were available from two high-quality trials.<sup>18,19</sup> At 4 months, Cunningham et al<sup>18</sup> reported a greater

increase in daily step count (1575; 95% CI 732 to 2419;  $p<0.001$ ). In contrast, Tew et al.,<sup>19</sup> did not find improvement in daily step count of their patients after a six-week intervention (440; 95% CI -827 to 1708;  $p>0.05$ ).

### ***Walking capacity***

Five studies<sup>18-20,22,23</sup> reported outcomes on walking capacity. One pretest-posttest study<sup>23</sup> reported significant improvement in ACD: increases of 63% at 3 months, and 84% at 6 months compared to baseline. Similarly, one high-quality pilot RCT<sup>19</sup> reported a significant increase in ACD at 6 weeks (173; 95% CI 23-332). Another high-quality trial<sup>20</sup> found no change in the number of people whose ACD improved, remained same or deteriorated at 2-month follow up. However, a significantly greater number of patients were found to have improved their maximum walking distance ( $p=0.003$ ) at 12 months.

### ***Quality of life and other outcomes***

Five trials,<sup>18-20,22,23</sup> reported outcomes related to quality of life (QoL). Generally, outcomes of both general and disease specific QoL were mixed. One high quality trial showed improvement in SF-36 (0.40 vs -0.30;  $p=0.002$ ) but no change in ICQ score ( $p>0.05$ ) at 4 months.<sup>18</sup> In contrast, another high quality pilot trial<sup>19</sup> reported improvement in ICQ (-10.6; 95% CI -18.9 to -2.3) but not general QoL (EQ-5D utility score 0.05; 95% CI -0.09 to 0.19). Two high-quality trials reported no change in health related QoL at two months (0.83 vs 0.85;  $p=0.33$ ), and at 12-months follow-up (0.83 vs 0.84;  $p=0.13$ )<sup>20</sup> or in general QoL at 14 weeks.<sup>22</sup> A low quality trial however, reported post-intervention improvement in the physical component of SF-36 score at both 3 months (40.8 vs 36.0;  $p<0.01$ ) and at 12-month follow-up (42.3 vs 36.0;  $p<0.01$ ) compared to baseline.<sup>33</sup> For this trial, there was also an improvement in the mental composite score at 3 months (45.2 vs 41.6;  $p<0.05$ ) but not at 12 months (44.2 vs 41.6;  $p>0.05$ ).

A high-quality trial reported an improved walking performance self-efficacy (mean adjusted difference 29.5; 95% CI 12.6-46.4) and perceived control over illness (mean adjusted difference 2.4 95% CI 0.0-4.7).<sup>19</sup> A low-quality trial<sup>23</sup> reported a decrease in pain intensity at 3 months compared to baseline (5.89 vs 4.73,  $p < 0.05$ ), which remained stable during the following 9 months (5.89 vs 4.53;  $p < 0.05$ ). Also, the time taken for claudication pain to disappear improved at 6 month from baseline (3.95 vs 2.01;  $p < 0.05$ ), and remained stable for the following 6 months (3.95 vs 2.83;  $p < 0.05$ ).

### ***Patients experiences with interventions***

Three studies<sup>18,19,23</sup> reported qualitative findings related to the experiences of patients with education interventions. Sixteen qualitative findings were extracted from the papers and grouped into eight categories. These categories were then merged to create three synthesised themes: Acquiring knowledge; Receiving pragmatic and tailored care; and Gaining confidence and self-monitoring (Tables 4 and 5). Although verbatim quotes were not available from the included papers, each synthesised theme is described below with some examples which reflect the patients' experiences, as reported in the included papers.

***Acquiring knowledge:*** Acquiring knowledge entails patients being provided information about their pathology and the systemic nature of IC, being informed of the importance of secondary prevention and risk factor modification including the importance of PA. Tew et al.<sup>19</sup> reported that patients valued a 6-week programme of pragmatic group-based structured patient education because it provided them with greater understanding of their condition, and empowered them to walk more.

***Receiving pragmatic and tailored care:*** Patients valued the idea of a group-based intervention which also gave opportunity for individual attention. Prevost et al.<sup>23</sup> reported that 97% of the participants were satisfied with the quality, duration, topics discussed as well as

the group nature of the session. Similarly, 95% reported that they were satisfied with the scope, quality, and benefit of individual discussion.

***Gaining confidence and self-monitoring:*** Gaining confidence encompassed developing a positive self-attitude, overcoming uncertainties, and feeling empowered. Gaining confidence to self-monitor their daily step counts with a pedometer: patients were presented with the opportunity to self-monitor their progress and they considered this an important component for meeting their physical activity goal. Prevost et al.<sup>23</sup> reported that their participants were very satisfied with the improvement in their attitude toward walking with claudication and their physical self-confidence and valued the use of pedometer as a tool for motivation, self-monitoring, and goal setting.

## **Discussion**

Six trials evaluating a range of patient education interventions for improving PA in patients with PAD and IC were reviewed. The included studies were mostly of high methodological quality. The major sources of risks of bias were lack of blinding of outcome assessment, performance bias and lack of allocation concealment. Combined evidence from four studies indicated that structured patient education intervention increased maximal walking capacity<sup>19,23</sup> and improved daily PA,<sup>18,20</sup> however, similar number of studies demonstrated no change in maximal walking capacity,<sup>20,22</sup> and free-living PA.<sup>19,20</sup> Generally, while daily step count tended to improve only after longer period of follow up (4 months upwards), there was no such trend for the outcomes related to maximal walking distance. Evidence from five trials<sup>18-20,22,23</sup> showed marked variability regarding quality of life measures. Useful interventions from the patients' perspective included those that provided them with information about their condition, were designed to enhance group interaction while maintaining individual discussion, and that provided them with confidence and self-monitoring.

Related reviews have been conducted on behaviour change techniques<sup>24</sup> and home-based exercise programmes<sup>25</sup> for PA improvement in this population. Galea et al.<sup>24</sup> reported limited evidence from one high-quality RCT to support BCTs for improving maximal and pain free walking capacities, and for increasing daily PA among people with IC. Similarly, Al-Jundi et al.<sup>25</sup> reported that there is “low-level” evidence that home-based exercise programmes can improve walking capacity and quality of life in patients with IC. However, the reviews were limited to RCTs of BCT,<sup>24</sup> or primarily reported outcomes related to walking capacity rather than daily PA.<sup>24,25</sup> No review has considered the qualitative experiences of patients with these interventions. Further, although these reviews include studies with patient education modalities, study eligibility did not specifically consider the key criteria for a structured patient education.<sup>13</sup>

Possible explanations for the contrasting findings in the current review may be related to the heterogeneity of study design. One<sup>23</sup> of the studies employed a prepost-posttest design therefore lacked a control group, and had high risk of bias including selection, detection, and performance bias. In another trial,<sup>21</sup> it was difficult to completely rule out possible contamination of the control group as both groups received some form of patient education. One study<sup>22</sup> included patients who underwent vascular intervention in the weeks prior to the intervention, and one patient who did not report claudication pain during treadmill testing. Furthermore, the large variations in patient education modalities and components warranted different contact time, duration of intervention, time point of outcome assessment, and intervention components, possibly resulting in wider outcome variability.

The high-quality pilot trial by Tew et al.,<sup>19</sup> which applied a 6-week structured education intervention demonstrated potential for increasing PA early in the programme, was particularly rigorous with blinded outcome assessment demonstrating improvement in walking capacity at 6 weeks. Although no change in daily step count was observed, it is

possible that the effect of behaviour change interventions takes longer to be noticeable. There could be other factors that mediate a slow response to adapting PA change in patients with PAD and IC even when the disease pathology is understood. One possible barrier is the claudication pain which these patients experience even when they are motivated to walk. Current NICE guidelines recommend “encouraging claudicants to exercise to the point of maximal pain.”<sup>7</sup> Perhaps for patients with IC to gain the benefit of secondary prevention, concomitant pain management may be desirable to delay the onset and reduce the intensity of pain. By delaying peak maximal pain and empowering patients through education the potential therapeutic value of walking not only could be realised early in the programme, but may be sustained.

Structured Exercise Programmes are the recommended exercise therapy for IC, however, based on the small number of included trials, most exercise programmes either do not incorporate patient education, or do not typically comply with criteria definitions of a structured education programme.<sup>12</sup> While ‘usual care’ patient education is included in the NICE SEPs recommendations for secondary preventions in IC, the development and implementation of the education components in SEPs interventions are not sufficiently reported to allow a judgement of their fidelity. Therefore, it is often not known how education was delivered, whether educators were sufficiently trained, whether education interventions were delivered as intended, and/or whether sufficient topics are included to enhance patient self-management skills. These elements are important as not only do they underpin the potential effectiveness of education programmes, they may be crucial in translating gains in a hospital based SEPs intervention to sustained improvement in daily physical activity at the end of the 3 months SEP duration, which is crucial in managing chronic diseases such as PAD/IC.

Only three of the included papers reported patient experience with interventions. In these trials, patients reported that they valued interventions that improved their understanding of the diseases pathology of PAD and IC, and provided them with information of the importance of walking and how walking helped. For these patients, gaining confidence, self-monitoring ability and skills were key to their accepting and perhaps adhering to interventions. Other important components according to the patients were the benefits from socialising in a group session without losing the opportunity for individualise care. The interventions that included these components demonstrated improvement in both maximal walking capacity,<sup>19,23</sup> and free-living PA.<sup>17</sup>

Several limitations are recognised regarding the conclusions in this review. First, the planned meta-analysis could not be implemented due to heterogeneity in the included studies.

Secondly, although four of the six included papers were of high quality, three were pilot trials assessing outcomes after a relatively short time point. For instance, Tew et al.<sup>19</sup> only had 13 or 9 participants in each group and assessed outcomes after just a 6-week intervention. This means that an inadequate sample and early assessment of outcome could limit the statistical power. Thirdly, the time point of outcome assessments were so varied that it was difficult to establish a reference time point to assess outcome performance even in a narrative synthesis.

Fourthly, lack of a control group or use of an active control group meant that the specific efficacy of the intervention versus the usual care or nothing could not be clearly ascertained.

In addition, the review design itself has inherent limitations; non-English language literature and unpublished literature was not searched or reviewed.

Limited evidence from four trials is inconclusive regarding the effect of structured patient education to improve daily PA, and walking capacity of patients with IC. Interventions that provided patients with information about their disease pathology and walking, provided motivation and empowered patients were valued, acceptable and seen by patients as



important in improving their PA. Structured education programmes may prove to be an essential part of exercise programmes for patients with IC but, rigorous trials are required before this can be recommended. Future interventions should consider the key criteria for a structured patient education programme. In addition, reporting on patients constructs of their experiences and perceptions to the interventions should always be factored in and reported so that holistic evaluation of effective components from the patients' perspective can be undertaken.

### **Acknowledgements**

No other persons besides the authors have made substantial contributions to this manuscript.

### **Source of funding**

UOA received funding from the University of Nigeria for a PhD at Glasgow Caledonian University, of which this work forms part.

**Conflict of interest:** none.

### **References**

1. McDermott MM. Lower extremity manifestations of peripheral artery disease: the pathophysiologic and functional implications of leg ischemia. *Circ Res.* 2015; 116:1540-50.
2. Doraiswamy VA, Giri J, Mohler E. Premature peripheral arterial disease – difficult diagnosis in very early presentation. *Int J Angiol.* 2009; 18(1): 45–47.
3. Mukherjee D, Cho L. Peripheral arterial disease: considerations in risks, diagnosis, and treatment. *J Natl Med Assoc.* 2009;101(10): 999-1008.
4. Booth FW, Roberts CK, Laye MJ. Lack of exercise is a major cause of chronic diseases. *Compr Physiol.* 2012;2(2):1143-1211. doi:10.1002/cphy.c110025.

5. Regensteiner JG, Hiatt WR, Coll JR, Criqui MH, TreatJacobson D, McDermott MM, et al. The impact of peripheral arterial disease on health-related quality of life in the Peripheral Arterial Disease Awareness, Risk, and Treatment: New Resources for Survival (PARTNERS) Program. *Vasc Med* 2008;13:15-24.
6. National Institute for Health and Care Excellence. NICE Peripheral arterial disease quality standard 2014. Assessed on 12<sup>th</sup> June 2016 from:<https://www.nice.org.uk/guidance/qs52/resources/peripheral-arterial-disease-20987238166458>
7. Fokkenrood HJ, Bendermacher BL, Lauret GJ, Willigendael EM, Prins MH, Teijink JA. Supervised exercise therapy versus nonsupervised exercise therapy for intermittent claudication. *Cochrane Database Syst Rev* 2013;8:CD005263.
8. Fokkenrood HJ, Lauret GJ, Verhofstad N et al. The effect of supervised exercise therapy on physical activity and ambulatory activities in patients with intermittent claudication. *Eur J Vasc Endovasc Surg*. 2015 49(2):184-91.
9. Garg PK, Tian L, Criqui MH et al Physical activity during daily life and mortality in patients with peripheral arterial disease. *Circ* 2006; 114:242-8.
10. Barbosa JP, Farah BQ, Chehuen MR, et al. Barriers to physical activity in patients with intermittent claudication. *Int J Behav Med* 2015; 22:70-6.
11. Gorely T, Crank H, Humphreys L, Nawaz S, Tew GA. Standing still in the street’’: Experiences, knowledge and beliefs of patients with intermittent claudication—A qualitative study. *J Vasc Nurs* 2015; 33(1):4-9.
12. Department of Health, Diabetes UK. Structured patient education in diabetes: Report from the Patient Education Working Group 2005. Accessed 10<sup>th</sup> October, 2016 at <https://www.diabetes.org.uk/Documents/Reports/StructuredPatientEd.pdf>.

13. Abaraogu UO, Dall PM, Seenan CA. Patient education interventions to improve physical activity in patients with intermittent claudication: A protocol for systematic review. *BMJ Open* 2016;6:e011405. doi:10.1136/bmjopen-2016-011405.
14. Cochrane Consumers and Communication Review Group. Communication review group: data extraction template for Cochrane reviews. Cochrane Collaboration, 2007.
15. Noyes J & Lewin S. Chapter 5: Extracting qualitative evidence. In: Noyes J, Booth A, Hannes K, Harden A, Harris J, Lewin S, Lockwood C (editors), *Supplementary Guidance for Inclusion of Qualitative Research in Cochrane Systematic Reviews of Interventions*. Version 1 (updated August 2011). Cochrane Collaboration Qualitative Methods Group, 2011. Available from URL <http://cqrmg.cochrane.org/supplemental-handbook-guidance>.
16. Higgins J, Green S. *Cochrane handbook for systematic reviews of interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. <http://www.cochrane-handbook.org>, 2014.
17. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015statement. *Syst Rev*. 2015; 4(1) doi: [10.1186/2046-4053-4-1](https://doi.org/10.1186/2046-4053-4-1)
18. Cunningham MA, Swanson V, O'Carroll RE, Holdsworth RJ. Randomized clinical trial of a brief psychological intervention to increase walking in patients with intermittent claudication. *Br J Surg*. 2012; 99(1):49-56. doi: 10.1002/bjs.7714.

19. Tew GA, Humphreys L, Crank H et al. The development and pilot randomised controlled trial of a group education programme for promoting walking in people with intermittent claudication. *Vasc Med*.2015; 20(4) 348–357.
20. Fowler B, Jamrozik K, Norman P, Allen Y, Wilkinson E. Improving maximum walking distance in early peripheral arterial disease: randomised controlled trial. *Aust J Physiother*. 2002; 48(4):269-75.
21. Collins TC, Krueger PN, Kroll TL, Sharf BF. Face-to-face interaction compared with video watching on use of physical activity in peripheral arterial disease: a pilot trial. *Angiol*. 2009; 60(1):21-30. doi: 10.1177/0003319708318382.
22. Mays RJ, Hiatt WR, Casserly IP, Rogers RK, Main DS, Kohrt WM, Ho PM, Regensteiner JG. Community-based walking exercise for peripheral artery disease: An exploratory pilot study. *Vasc Med*. 2015; 20(4):339-47. doi: 10.1177/1358863X15572725.
23. Prévost A, Lafitte M, Pucheu Y, Couffinhal T; on behalf the CEPTA educational team. Education and home based training for intermittent claudication: functional effects and quality of life. *Eur J Prev Cardiol*. 2015; 22 (3):373-9. doi: 10.1177/2047487313512217.
24. Galea MN, Weinman JA, White C, Bearne L M. Do Behaviour-Change Techniques Contribute to the Effectiveness of Exercise Therapy in Patients with Intermittent Claudication? A Systematic Review. *Eur J Vasc Endovasc Surg*. 2013; 46:132-141.
25. Al-Jundi W, Madbak K, Beard JD, Nawaz S, Tew GA. Systematic Review of Home-based Exercise Programmes for Individuals with Intermittent Claudication. *Eur J Vasc Endovasc Surg*. 2013; 46(6): 690.




Figure 1: Patient Education review PRISMA Flow Diagram

Table 1: Characteristics of included studies

Study (country), Design, Quality, Attrition	Participants	Descriptions of Interventions	Outcomes, follow-up	Conclusion
Cunningham et al. <sup>18</sup> (UK)  RCT  High quality  3%	Total $n = 58$ , IC in at least one leg Post-exercise ABI<0.7	Intervention: Usual care + Information on PAD and walking, motivational interviewing, PA goal setting, action planning, self-monitoring and feedback, barrier identification with problem solving. Delivered at patients' homes by via a trainee health psychologist 2 x 1-hr sessions  Control: Usual care (walking advice + consultation with vascular surgeon)	Daily steps (pedometer)  Walking ability (patient report)  ICQ (disease-specific quality of life)  Medical Outcome Study Short-Form (generic health-related quality of life)  Outcomes assessed at baseline and 4 months	Brief psychological intervention significantly improved walking behaviour in patients with IC in comparison with usual care.
Tew et al. <sup>19</sup> (UK)  Pilot RCT  High quality  4.35%	Total $n = 23$  Stable Rutherford IC Classification 1-3 for $\geq 3$ months.	Intervention: Usual care plus one off 3-hr session of group patient centred structured education including patient story, PAD/IC and walking information provision, PA feedback, barrier identification with problem solving, goal setting, action planning, and self-monitoring. Delivered at in clinical research facility followed by twice-weekly phone calls for 6 weeks.  Control: Usual care	Daily step (accelerometer)  ICD, 6-minute walk test  WELCH questionnaire  ICQ (disease-specific quality of life)  EQ-5D (generic health-related quality of life)  Self-efficacy  Acceptability (exit	Education programme is feasible, acceptable, and potentially useful for improving walking capacity and quality of life

			interview)	
			Outcomes assessed at baseline and 6 weeks	
Fowler et al. <sup>20</sup> (Australia)	Total $n = 882$	Intervention: Educational package + mobility program (supervised or home-based) + smoking cessation where applicable	ICD	Intervention for early PAD based on increased PA & Smoking cessation results in a greater max walking distance 12 months later.
RCT	ABI $\leq 0.9$	Control: No intervention	PA pattern (Patient report)	
High quality	Clinical diagnosis of PAD Definite IC or atypical IC on Edinburgh Claudication Questionnaire		Medical Outcome Study Short-Form (generic health-related quality of life)	
15%			Outcomes assessed at baseline, 2 months, and 12 months	
Collins et al. <sup>21</sup> (USA)	Total $n = 51$ , Diagnosis of PAD ABI $> 0.5$ & $\leq 0.955$ .	Intervention: Communication intervention beginning with completion of guide and followed by 15-20 minutes of motivational interviewing, provision of information related to PAD/IC and walking, barrier identification and problem solving strategies, patient tailored walking prescription. Delivered by medical student.	PA pattern and time (patients report)	-Patients watching video on the use of PA in PAD improved participants walking speed
Pilot RCT			ICD and ACD	
Low quality			WIQ	
13.7%		Control: Video Intervention	Outcomes assessed at baseline, and 12 weeks.	
Prevost et al. <sup>23</sup> (France)	Total $n = 48$	Intervention: Educational classes, implementation of secondary prevention, and a personalised program of reconditioning exercises.	ICD and ACD	Educational therapeutic program results in a significant improvement in functional and QoL parameters during the first 3 months in patients with IC and persists even patients are no longer coached.
Pretest-posttest design	Level II Leriche and Fontaine IC Atypical symptom ABI $< 0.9$ ;	Control: No intervention	Medical Outcome Study Short-Form (generic health-related quality of life)	
Low quality				

4.17%			Pain intensity	
			Patients experiences via exit questionnaire (12months only)	
			Outcomes assessed at baseline, 3 months and 12 months.	
Mays et al. <sup>22</sup> (USA)	Total <i>n</i> = 25	Intervention: Initial in-hospital walking exercise (2 weeks, 3 days/week) followed by community-based walking exercise (12 weeks) with training, monitoring and coaching (TMC) components.	ICD and ACD	Community-based walking exercise with TMC improves ICD and walking performances other than ACD.
Pilot RCT	Patients without walking limiting comorbidities except IC		WIQ	
High quality	Severe cardiac ischemia ≤3months previous myocardia infarction TIA or stroke	Control: Usual care (standard advice to walk)	Medical Outcome Study Short-Form (generic health-related quality of life)	
20%	< 1month treatment with cilostazol or pentoxifylline Endovascular therapy 4-6wks prior baseline Or stable IC without revascularisation in the last 4-6weeks (ABI ≤0.9)		Outcomes assessed at baseline and 14 weeks	
Key: IC = intermittent claudication; ABI = Ankle brachial index; RCT; Randomised control trial; ACD = Absolute claudication distance; ICD = Initial claudication distance; NS = not significant, PA= Physical activity, WIQ =walking impairment questionnaire; ICQ = Intermittent claudication questionnaire, EQ-5D = EuroQoL EQ-5D-5L, WELCH = Walking Estimated- Limitation Calculated by History questionnaire				



Table 2: Risk of bias in individual studies

Authors	*Sources/ Potential sources of bias						Summary of risk of bias	†Quality index	
	Selection bias		Performance bias	Detection bias	Bias due to attrition	Reporting bias			Other bias
	Random sequence generation	Allocation concealment	Participants and personnel blinding	Blinding of outcome assessments	Incomplete outcome data	Selective reporting			
Cunningham et al. <sup>18</sup>	No	Yes	No	Yes	No	No	Trial was not powered	Low	High
Tew et al. <sup>19</sup>	No	No	No	No	Yes	No	Pilot study	Low	High
Fowler et al. <sup>20</sup>	No	No	No	Yes	No	No	NA	Low	High
Collins et al. <sup>21</sup>	No	Yes	Yes	Yes	No	No	Pilot study	High	Low
Prevost et al. <sup>23</sup>	Yes	Yes	Yes	Yes	No	No	NA	High	Low
Mays et al. <sup>22</sup>	No	No	Yes	Yes	No	No	Pilot study	Low	High

<sup>#</sup>The Cochrane Collaboration tool for assessing risk of bias was used to determine and summarise possible sources of risk of bias in included studies (Cochrane 2011) (\*Yes indicates the presence or potential presence of a source of bias). <sup>\*</sup>Summary risk of bias in included studies was presented. <sup>†</sup>Studies were subsequently rated as low-quality trials (i.e. having high risk of bias) or high-quality trials (i.e. having low to moderate risk of bias) if there was  $\geq 3$  or  $< 3$  identifiable sources of bias respectively

Table 3: Data extraction of finding from included studies (Except where specified, results are presented as intervention group compared to control group)

Authors	Change in Daily PA behaviour	PA Capacity/Ability measures	Pain, Self-efficacy, and Perceived control over illness.	Quality of life
Mays et al. <sup>22</sup>		At 14 weeks: Pain free walking time (p= NS);  Greater increase in caudication onset time (5.8±1.5min to 7.4±1.6min vs 4.7±1.4min → 4.1±1.5min; p=0.045);  Greater improvement in walking impairment (42.3±7.7min → 60.6±7.2% vs 49.1±7.7 → 44.6±7.2%; p=0.001).		At 14 weeks:  General QoL (p=NS)
Prevost et al. <sup>23</sup>		Significant % increase in ICD from baseline (277% at 3months, 203% @ 6months, 141% @ 12months; p<0.001)  Significant % increase in ACD from baseline (63% @ 3months; 84% @ 6months; 65% @ 12month; p <0.01).	Decrease in pain_intensity at 3, 6, 12months from baseline (5.89→ 4.73* → 4.34 <sup>a</sup> → 4.53*)  Improved from baseline in Time of release of pain at 6 months, and 12months(3.95 → 2.01 <sup>a</sup> → 2.83 <sup>a</sup> )	Improvement from baseline in physical composite score of SF-36 at 3, 6, & 12months (36.0→ 40.8→ 41.9→ 42.9; p<0.01)  Improvement from baseline in the mental composite of SF-36 at 3 and 6 months (41.6→ 45.5→ 44.7→ 44.2; p<0.05)
Cunningham et al. <sup>18</sup>	At 4 months: Greater increase in daily steps (1358 vs -227; p<0.001)	At 4 months: *Greater pain-free walking distance (1.00 vs 0.00; p=0.008)		At 4 months: Improvement in general QoL (0.40 vs -0.30; p=0.002)  Disease specific QoL (p=NS)
Tew et al., 2015	At six weeks : Daily steps (p=NS)	At six weeks: Improvement in Six-minute walk distance (44.9; CI 6.9 to 82.9); Greater increase in ACD (173; CI 23 to 322); ICD p=NS Greater improvement in self-reported walking ability WELCH score (21.8; CI 8.6 to 35.0); WIQ Speed (21.0; CI 3.8 to 38.1), WIQ distance (30.7; CI 6.4 to 55.0), WIQ stair climbing (30.7; CI 6.4 to 55.0).	At six weeks: Improvement in walking performance efficacy (29.5; CI 12.6 to 46.4)  Greater improvement in perceive control over illness (2.4; CI 0.0 to 4.7)	At six weeks: Greater improvement in disease specific QoL (-10.6; CI -18.9 to -2.3)  General QoL (p=NS)
Fowler et	At 2nd months: % of patients walking for	At 2nd month: Self-report maximum		At 2nd month: HQoL(NS)

al., 2002	recreation $\geq 3$ /wk ( $p=NS$ ); % of patients engaging in vigorous PA ( $p=NS$ ); % of patient belonging to exercise group ( $p=NS$ )  <b>At 12th month:</b> Greater % of patient walking for recreation $\geq 3$ /wk (33.8 vs 25; $p=0.01$ ); % of patients engaging in vigorous PA ( $p=NS$ ); Greater % of belonging to exercise group (16.5 vs 1.8, $p<0.001$ )	walking distance before the onset of pain ( $p=NS$ )  <b>At 12th month:</b> Improvement in self-report maximum walking distance before the onset of pain ( $p=0.04$ )		At 12th month HQoL(NS)
Collins et al 2009	At 12weeks: Activity patterns in various levels of physical activity ( $p=NS$ )			

Keys: PWD = Peak walking distance; WIQ = Walking impairment questionnaire; QoL = Quality of life; COT = Claudication onset time; Int = Intervention; Cont = Control; grp = Group; NS = No significance;  $\Delta$  = Change; HQoL = Health related quality of life; MWD = Maximum walking distance; ACD = Absolute claudication distance; PA = Physical activity; ICD = Initial claudication distance; FCD = Functional claudication distance.

\*Results adjusted for co-founding variables

<sup>#</sup> $p<0.05$ , <sup>u</sup> $p<0.01$

Table 4: Categories: Label and description of patients experiences with intervention

Category label	Category description
Receiving information about disease	Participants valued participation in the intervention because it provided them with greater understanding of their condition (T); Patient valued intervention because it provided them with extra information about their illness(C). 97% were very satisfied with the topic discussed (P)
Receiving information about walking	Patients reported that they valued the intervention because it provided them with understanding of the importance of walking(C); Patients reported that they valued the intervention because it provided them with understanding of how walking will help(C); Patient said intervention were worthwhile and that they valued it because it provided them with extra information about walking(C)
Being motivation and empowered	Patients reported being satisfied with their improvement in attitude towards walking with their claudication and their physical self-confidence (P); Patients reported being satisfied with their improvement in their physical self-confidence (P); Participants valued participation because it enabled them to walk more (T); Patients said they valued intervention because it provided them with extra encouragement and motivation(C).
Benefit of group education session	97% reported that they were very satisfied with the benefit of group education session (P)
Self-monitoring	The use of pedometer was valued as it was seen as useful tool for self-monitoring (T)
Goal setting	Patient reported that intervention were worthwhile because it provided them clarity on what to do(C);
Pedometer as useful tool	Patients valued the pedometer and seen as a valuable tool for motivation, self-monitoring and goal setting (T)
Receiving personalised care	95.5% reported very satisfied with the scope, quality, and benefit of individual discussion(P); Patients valued the intervention because it provided personalised plan(C)
Key: T: Tew et al. <sup>19</sup> ; C: Cunningham et al. <sup>18</sup> ; P: Prevost et al. <sup>23</sup>	

Table 5: Developing synthesis findings from the categories

Synthesised finding	Category
Acquiring knowledge	Receiving information about disease; Receiving information about for walking; Goal setting; Pedometer as useful tool
Pragmatic and tailored care	Benefit of group education session; Receiving personalised care
Gaining confidence and self-monitoring	Being motivation and empowered; Pedometer as useful tool; Self-monitoring; Pedometer as useful tool